


LISTING OF THE CLAIMS

Claim 1 (currently amended): A method of treating cytokine-mediated cancer said method comprising administering to a patient in need of such treatment a therapeutically effective amount of a compound of the formula (I):



wherein

Ar₁ is a pyrazole wherein Ar₁ may be substituted by one or more R₁, R₂ or R₃;

Ar₂ is:

phenyl, naphthyl, quinoline, isoquinoline, tetrahydronaphthyl, tetrahydroquinoline, tetrahydroisoquinoline, benzimidazole, benzofuran, indanyl, indenyl or indole each being optionally substituted with one to three R₂ groups;

L is a C₁₋₁₀ saturated or unsaturated branched or unbranched carbon chain;

wherein one or more methylene groups are optionally independently replaced by O, N or S; and

wherein said linking group is optionally substituted with 0-2 oxo groups and one or more C₁₋₄ branched or unbranched alkyl which may be substituted by one or more halogen atoms;

Q is selected from the group consisting of:

a) pyridine, pyrimidine, pyridazine, imidazole, benzimidazole, oxazo[4,5-*b*]pyridine and imidazo[4,5-*b*]pyridine, which are optionally substituted with one to three groups selected from

the group consisting of halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, hydroxy, mono- or di-(C₁₋₃ alkyl)amino, C₁₋₆ alkyl-S(O)_m and phenylamino wherein the phenyl ring is optionally substituted with one to two groups consisting of halogen, C₁₋₆ alkyl and C₁₋₆ alkoxy;

b) morpholine, thiomorpholine, thiomorpholine sulfoxide, thiomorpholine sulfone, piperidine, piperidinone and tetrahydropyrimidone which are optionally substituted with one to three groups selected from the group consisting of C₁₋₆ alkyl, C₁₋₆ alkoxy, hydroxy, mono- or di-(C₁₋₃ alkyl)amino-C₁₋₃ alkyl, phenylamino-C₁₋₃ alkyl and C₁₋₃ alkoxy-C₁₋₃ alkyl;

R₁ is selected from the group consisting of:

- a) C₃₋₁₀ branched or unbranched alkyl, which may optionally be partially or fully halogenated, and optionally substituted with one to three phenyl, naphthyl or heterocyclic groups selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, isoxazolyl and isothiazolyl; each such phenyl, naphthyl or heterocycle selected from the group hereinabove described, being substituted with 0 to 5 groups selected from the group consisting of halogen, C₁₋₆ branched or unbranched alkyl which is optionally partially or fully halogenated, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkenyl, hydroxy, cyano, C₁₋₃ alkyloxy which is optionally partially or fully halogenated, NH₂C(O) and di(C₁₋₃)alkylaminocarbonyl;
- b) C₃₋₇ cycloalkyl selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl and bicycloheptanyl, which may optionally be partially or fully halogenated and which may optionally be substituted with one to three C₁₋₃ alkyl groups, or an analog of such cycloalkyl group wherein one to three ring methylene groups are replaced by groups independently selected from O, S, CHOH, >C=O, >C=S and NH;
- c) C₃₋₁₀ branched alkenyl which may optionally be partially or fully halogenated, and which is optionally substituted with one to three C₁₋₅ branched or unbranched alkyl, phenyl, naphthyl or heterocyclic groups, with each such heterocyclic group being independently selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, isoxazolyl and isothiazolyl, and each such phenyl, naphthyl or heterocyclic group being substituted with 0 to 5 groups selected from halogen, C₁₋₆ branched or

unbranched alkyl which is optionally partially or fully halogenated, cyclopropyl, cyclobutyl, cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl and bicycloheptanyl, hydroxy, cyano, C_{1-3} alkyloxy which is optionally partially or fully halogenated, $NH_2C(O)$, mono- or di(C_{1-3})alkylaminocarbonyl;

d) C_{5-7} cycloalkenyl selected from the group consisting of cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptenyl, cycloheptadienyl, bicyclohexenyl and bicycloheptenyl, wherein such cycloalkenyl group may optionally be substituted with one to three C_{1-3} alkyl groups;

e) cyano; and,

f) methoxycarbonyl, ethoxycarbonyl and propoxycarbonyl;

R_2 is selected from the group consisting of:

a C_{1-6} branched or unbranched alkyl which may optionally be partially or fully halogenated, acetyl, aroyl, C_{1-4} branched or unbranched alkoxy, which may optionally be partially or fully halogenated, halogen, methoxycarbonyl and phenylsulfonyl;

R_3 is selected from the group consisting of:

a) a phenyl, naphthyl or heterocyclic group selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, tetrahydrofuryl, isoxazolyl, isothiazolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, benzoxazolyl, benzisoxazolyl, benzpyrazolyl, benzothiofuranyl, cinnolyl, pterindinyl, phthalazinyl, naphthypyridinyl, quinoxalyl, quinazolinyl, purinyl and indazolyl; wherein such phenyl, naphthyl or heterocyclic group is optionally substituted with one to five groups selected from the group consisting of a C_{1-6} branched or unbranched alkyl, phenyl, naphthyl, heterocycle selected from the group hereinabove described, C_{1-6} branched or unbranched alkyl which is optionally partially or fully halogenated, cyclopropyl, cyclobutyl, cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl, bicycloheptanyl, phenyl C_{1-5} alkyl, naphthyl C_{1-5} alkyl, halo, hydroxy, cyano, C_{1-3} alkyloxy which may optionally

be partially or fully halogenated, phenyloxy, naphthyloxy, heteraryloxy wherein the heterocyclic moiety is selected from the group hereinabove described, nitro, amino, mono- or di-(C₁₋₃)alkylamino, phenylamino, naphthylamino, heterocyclylamino wherein the heterocyclyl moiety is selected from the group hereinabove described, NH₂C(O), a mono- or di-(C₁₋₃)alkyl aminocarbonyl, C₁₋₅ alkyl-C(O)-C₁₋₄ alkyl, amino-C₁₋₅ alkyl, mono- or di-(C₁₋₃)alkylamino-C₁₋₅ alkyl, amino-S(O)₂, di-(C₁₋₃)alkylamino-S(O)₂, R₄-C₁₋₅ alkyl, R₅-C₁₋₅ alkoxy, R₆-C(O)-C₁₋₅ alkyl and R₇-C₁₋₅ alkyl(R₈)N;

b) a fused aryl selected from the group consisting of benzocyclobutanyl, indanyl, indenyl, dihydronaphthyl, tetrahydronaphthyl, benzocycloheptanyl and benzocycloheptenyl, or a fused heterocyclyl selected from the group consisting of cyclopentenopyridine, cyclohexanopyridine, cyclopentanopyrimidine, cyclohexanopyrimidine, cyclopentanopyrazine, cyclohexanopyrazine, cyclopentanopyridazine, cyclohexanopyridazine, cyclopentanoquinoline, cyclohexanoquinoline, cyclopentanoisoquinoline, cyclohexanoisoquinoline, cyclopentanoindole, cyclohexanoindole, cyclopentanobenzimidazole, cyclohexanobenzimidazole, cyclopentanobenzoxazole, cyclohexanobenzoxazole, cyclopentanoimidazole, cyclohexanoimidazole, cyclopentanothiophene and cyclohexanothiophene; wherein the fused aryl or fused heterocyclyl ring is substituted with 0 to 3 groups independently selected from phenyl, naphthyl and heterocyclyl selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, isoxazolyl, and isothiazolyl, C₁₋₆ branched or unbranched alkyl which is optionally partially or fully halogenated, halo, cyano, C₁₋₃ alkyloxy which is optionally partially or fully halogenated, phenyloxy, naphthyloxy, heterocycliloxy wherein the heterocyclyl moiety is selected from the group hereinabove described, nitro, amino, mono- or di-(C₁₋₃)alkylamino, phenylamino, naphthylamino, heterocyclylamino wherein the heterocyclyl moiety is selected from the group hereinabove described, NH₂C(O), a mono- or di-(C₁₋₃)alkyl aminocarbonyl, C₁₋₄ alkyl-OC(O), C₁₋₅ alkyl-C(O)-C₁₋₄ branched or unbranched alkyl, an amino-C₁₋₅ alkyl, mono- or di-(C₁₋₃)alkylamino-C₁₋₅ alkyl, R₉-C₁₋₅ alkyl, R₁₀-C₁₋₅ alkoxy, R₁₁-C(O)-C₁₋₅ alkyl and R₁₂-C₁₋₅ alkyl(R₁₃)N;

c) cycloalkyl selected from the group consisting of cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl and bicycloheptanyl, which the cycloalkyl may

optionally be partially or fully halogenated and which may optionally be substituted with one to three C_{1-3} alkyl groups;

d) C_{5-7} cycloalkenyl, selected from the group consisting of cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptenyl, cycloheptadienyl, bicyclohexenyl and bicycloheptenyl, wherein such cycloalkenyl group may optionally be substituted with one to three C_{1-3} alkyl groups; and

e) acetyl, aroyl, alkoxycarbonylalkyl or phenylsulfonyl;

f) C_{1-6} branched or unbranched alkyl which may optionally be partially or fully halogenated;

wherein

or R_1 and R_2 taken together may optionally form a fused phenyl or pyridinyl ring,

each R_8 , R_{13} is independently selected from the group consisting of:

hydrogen and C_{1-4} branched or unbranched alkyl which may optionally be partially or fully halogenated;

each R_4 , R_5 , R_6 , R_7 , R_9 , R_{10} , R_{11} and R_{12} is independently selected from the group consisting of:

morpholine, piperidine, piperazine, imidazole and tetrazole;

$m = 0, 1, 2$;

$r = 0, 1, 2$;

$t = 0, 1, 2$;

and

$X = O$ or S or

the physiologically acceptable acids or salts thereof.

Claim 2 (original) The method according to claim 1 wherein Ar₂ is naphthyl, tetrahydronaphthyl, indanyl or indenyl.

Claim 3 (original) The method according to claim 2 wherein Ar₂ is naphthyl.

Claim 4 (previously presented): The method according to claim 3 wherein:

Ar₂ is 1-naphthyl;

L is C₁₋₆ saturated or unsaturated branched or unbranched carbon chain wherein one or more methylene groups are optionally independently replaced by O, N or S; and wherein said linking group is optionally substituted with 0-2 oxo groups and one or more C₁₋₄ branched or unbranched alkyl which may be substituted by one or more halogen atoms;

R₁ is selected from the group consisting of C₃₋₁₀ alkyl branched or unbranched, cyclopropyl and cyclohexyl which may optionally be partially or fully halogenated and which may optionally be substituted with one to three C₁₋₃ alkyl groups;

R₃ is selected from the group consisting of C₁₋₄ alkyl branched or unbranched, cyclopropyl, cyclopentyl, phenyl, pyridinyl each being optionally substituted as described in claim 1 and alkoxy carbonyl alkyl.

Claim 5 (canceled)

Claim 6 (currently amended): The method according to claim 5 wherein L is C₁₋₅ saturated carbon chain wherein one or more methylene groups are optionally independently replaced by O, N or S; wherein said linking group is optionally substituted with 0-2 oxo groups and one or more C₁₋₄ branched or unbranched alkyl which may be substituted by one or more halogen atoms; and X = O.

Claim 7 (original): The method according to claim 6 wherein L is propoxy, ethoxy or methoxy each being optionally substituted with 0-2 oxo groups and one or more C₁₋₄ branched or unbranched alkyl which may be substituted by one or more halogen atoms.

Claim 8 (original): The method according to claim 7 wherein L is ethoxy optionally substituted with 0-2 oxo groups and one or more C₁₋₄ branched or unbranched alkyl which may be substituted by one or more halogen atoms.

Claim 9 (original): The method according to claim 6 wherein L is methyl or propyl each being optionally substituted with 0-2 oxo groups and one or more C₁₋₄ branched or unbranched alkyl which may be substituted by one or more halogen atoms.

Claim 10 (original): The method according to claim 6 wherein L is C₃₋₅ acetylene optionally substituted with 0-2 oxo groups and one or more C₁₋₄ branched or unbranched alkyl which may be substituted by one or more halogen atoms.

Claim 11 (original): The method according to claim 6 wherein L is methylamino optionally substituted with 0-2 oxo groups and one or more C₁₋₄ branched or unbranched alkyl which may be substituted by one or more halogen atoms.

Claim 12 (previously presented): The method according to claim 1 wherein the compound is chosen from:

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-(*cis*-2,6-dimethylmorpholin-4-yl)ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-(*trans*-2,6-dimethylmorpholin-4-yl)ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-(2-(methoxymethyl)morpholin-4-yl)ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-(morpholin-4-yl)-2-oxoethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-(morpholin-4-yl)-2-methylethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-(morpholin-4-yl)-1-methylethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-thiomorpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-(1-oxothiomorpholin-4-yl)ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)-3-methylnaphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-(morpholin-4-yl-carbonyloxo)ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(3-morpholin-4-yl-propyl)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(morpholin-4-yl-methyl)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-pyridin-4-yl-ethyl)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(3-(morpholin-4-yl)propyn-1-yl)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(3-(piperidin-1-yl)propyn-1-yl)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(3-(2-methoxymethylmorpholin-4-yl)propyn-1-yl)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(pyridin-4-yl-methoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-pyridin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(3-pyridin-4-yl-propoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-imidazol-1-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(pyridin-4-yl-methylamino)naphthalen-1-yl]-urea;

1-[5-*iso*-Propyl-2-phenyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-cyclohexyl-2-phenyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-(2,2,2-trifluoroethyl)-2-phenyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-(1-methylcycloprop-1-yl)-2-phenyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-(1-methylcyclohex-1-yl)-2-phenyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-methyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(4-chlorophenyl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-butyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(4-methyl-3-carbamylphenyl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(4-methyl-3-(morpholin-4-yl)methylphenyl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(4-methyl-3-dimethylaminomethylphenyl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(3-dimethylaminomethylphenyl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(2-chloropyridin-5-yl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(2-methylpyridin-5-yl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(2-methoxypyridin-5-yl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(pyridin-3-yl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(2-methylpyridin-5-yl)-2H-pyrazol-3-yl]-3-[4-(2-pyridin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(2-methylpyridin-5-yl)-2H-pyrazol-3-yl]-3-[4-(2-(*trans*-2,6-dimethylmorpholin-4-yl)ethoxy)naphthalen-1-yl]-urea and

1-[5-*tert*-butyl-2-(2-methylpyridin-5-yl)-2H-pyrazol-3-yl]-3-[4-(3-morpholin-4-yl-propyn-1-yl)naphthalen-1-yl]-urea

or the physiologically acceptable acids or salts thereof.

Claim 13 (original): The method according to claim 12 wherein the compound is chosen from the group consisting of:

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-(1-oxothiomorpholin-4-yl)ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(2-methylpyridin-5-yl)-2H-pyrazol-3-yl]-3-[4-(2-pyridin-4-yl-ethoxy)naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(2-methoxypyridin-5-yl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea and

1-[5-*tert*-butyl-2-methyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea
or the physiologically acceptable acids or salts thereof.

Claim 14 (original): The method according to claim 13 wherein the compound is:

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea
or the physiologically acceptable acids or salts thereof.

Claim 15 (previously presented): The method according to claim 1 wherein the treatment is done in conjunction with genotoxic therapy.

Claim 16 (previously presented): A method of treating cytokine-mediated cancer said method comprising administering to a patient in need of such treatment a therapeutically effective amount of a compound chosen from

1-[5-(2-hydroxy-1,1-dimethyl-ethyl)-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)-naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(3-hydroxy-4-methyl-phenyl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)-naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)-naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-{4-[2-(3-oxo-morpholin-4-yl)-ethoxy]-naphthalen-1-yl}-urea;

1-[5-*tert*-butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-{4-[2-(4-oxy-morpholin-4-yl)-ethoxy]-naphthalen-1-yl}-urea;

1-[5-(2-hydroxy-1,1-dimethyl-ethyl)-2-(6-methyl-pyridin-3-yl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)-naphthalen-1-yl]-urea;

1-[5-*tert*-butyl)-2-(1-oxy-6-methyl-pyridin-3-yl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)-naphthalen-1-yl]-urea;

1-[5-*tert*-butyl)-2-(6-methyl-pyridin-3-yl)-2H-pyrazol-3-yl]- 3-{4-[2-(4-oxy-morpholin-4-yl)-ethoxy]-naphthalen-1-yl}-urea;

1-[5-(2-hydroxy-1,1-dimethyl-ethyl)-2-(6-methyl-pyridin-3-yl)-2H-pyrazol-3-yl]-3-[4-(2-pyridin-4-yl-ethoxy)-naphthalen-1-yl]-urea;

1-[5-*tert*-butyl)-2-(6-methyl-pyridin-3-yl)-2H-pyrazol-3-yl]-3-[4-(2-hydroxy-2-pyridin-4-yl-ethoxy)-naphthalen-1-yl]-urea;

1-[5-*tert*-butyl)-2-(6-methyl-pyridin-3-yl)-2H-pyrazol-3-yl]-3-{4-[2-(1-oxy-pyridin-4-yl)-ethoxy]-naphthalen-1-yl}-urea;

1-[5-(2-hydroxy-1,1-dimethyl-ethyl)-2-*p*-tolyl-2H-pyrazol-3-yl]- 3-{4-[2-(1-oxo-thiomorpholin-4-yl)-ethoxy]-naphthalen-1-yl}-urea;

1-[5-*tert*-butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]- 3-{4-[2-(1-oxo-thiomorpholin-4-yl)-ethoxy]-naphthalen-1-yl}-urea;

1-[5-*tert*-butyl-2-*p*-tolyl-2H-pyrazol-3-yl]- 3-{4-[2-(1,3 dioxo-thiomorpholin-4-yl)-ethoxy]-naphthalen-1-yl}-urea;

1-[5-(2-hydroxy-1,1-dimethyl-ethyl)-2-methyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)-naphthalen-1-yl]-urea;

1-[5-*tert*-butyl-2-methyl-2H-pyrazol-3-yl]-3-{4-[2-(4-oxy-morpholin-4-yl)-ethoxy]-naphthalen-1-yl}-urea;

1-[5-*tert*-Butyl-2-(2-hydroxy-4-methyl-phenyl)-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)-naphthalen-1-yl]-urea;

4-(3-*tert*-Butyl-5-{3-[4-(2-morpholin-4-yl-ethoxy)-naphthalen-1-yl]-ureido}-pyrazol-1-yl)-benzoic acid;

1-[5-(1,1-Dimethyl-2-oxo-ethyl)-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)-naphthalen-1-yl]-urea;

2-Methyl-2-(5-{3-[4-(2-morpholin-4-yl-ethoxy)-naphthalen-1-yl]-ureido}-1-*p*-tolyl-1H-pyrazol-3-yl)-propionic acid;

1-(5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl)-3-[4-(2-morpholin-4-yl-2-oxo-ethoxy)-naphthalen-1-yl]-urea and

1-(5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl)-3-{4-[2-(1-oxo-1 λ^4 -thiomorpholin-4-yl)-ethoxy]-naphthalen-1-yl}-urea


or physiologically acceptable acids or salts thereof.

Claim 17 (previously presented): The method according to claim 1 wherein the cancer is chosen from cancer cachexia, multiple myeloma and acute myelogenous leukemia.

Claim 18 (previously presented): The method according to claim 1 wherein the cancer is multiple myeloma.

Claim 19 (previously presented): The method according to claim 14 wherein the cancer is multiple myeloma.

Claim 20 (currently amended): A method of treating cytokine-mediated tumors said method comprising administering to a patient in need of such treatment a therapeutically effective amount of a compound of the formula (I):



wherein

Ar₁ is a pyrazole wherein Ar₁ may be substituted by one or more R₁, R₂ or R₃;

Ar₂ is:

phenyl, naphthyl, quinoline, isoquinoline, tetrahydronaphthyl, tetrahydroquinoline, tetrahydroisoquinoline, benzimidazole, benzofuran, indanyl, indenyl or indole each being optionally substituted with one to three R₂ groups;

L is a C₁₋₁₀ saturated or unsaturated branched or unbranched carbon chain;
wherein one or more methylene groups are optionally independently replaced by O, N or S; and

wherein said linking group is optionally substituted with 0-2 oxo groups and one or more C₁₋₄ branched or unbranched alkyl which may be substituted by one or more halogen atoms;

Q is selected from the group consisting of:

- a) pyridine, pyrimidine, pyridazine, imidazole, benzimidazole, oxazo[4,5-*b*]pyridine and imidazo[4,5-*b*]pyridine, which are optionally substituted with one to three groups selected from the group consisting of halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, hydroxy, mono- or di-(C₁₋₃ alkyl)amino, C₁₋₆ alkyl-S(O)_m and phenylamino wherein the phenyl ring is optionally substituted with one to two groups consisting of halogen, C₁₋₆ alkyl and C₁₋₆ alkoxy;
- b) morpholine, thiomorpholine, thiomorpholine sulfoxide, thiomorpholine sulfone, piperidine, piperidinone and tetrahydropyrimidone which are optionally substituted with one to three groups selected from the group consisting of C₁₋₆ alkyl, C₁₋₆ alkoxy, hydroxy, mono- or di-(C₁₋₃ alkyl)amino-C₁₋₃ alkyl, phenylamino-C₁₋₃ alkyl and C₁₋₃ alkoxy-C₁₋₃ alkyl;

R₁ is selected from the group consisting of:

- a) C₃₋₁₀ branched or unbranched alkyl, which may optionally be partially or fully halogenated, and optionally substituted with one to three phenyl, naphthyl or heterocyclic groups selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, isoxazolyl and isothiazolyl; each such phenyl, naphthyl or heterocycle selected from the group hereinabove described, being substituted with 0 to 5 groups selected from the group consisting of halogen, C₁₋₆ branched or unbranched alkyl which is optionally partially or fully halogenated, C₃₋₈ cycloalkyl, C₅₋₈ cycloalkenyl, hydroxy, cyano, C₁₋₃ alkyloxy which is optionally partially or fully halogenated, NH₂C(O) and di(C₁₋₃)alkylaminocarbonyl;

- b) C₃₋₇ cycloalkyl selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl and bicycloheptanyl, which may optionally be partially or fully halogenated and which may optionally be substituted with one to three C₁₋₃ alkyl groups, or an analog of such cycloalkyl group wherein one to three ring methylene groups are replaced by groups independently selected from O, S, CHOH, >C=O, >C=S and NH;
- c) C₃₋₁₀ branched alkenyl which may optionally be partially or fully halogenated, and which is optionally substituted with one to three C₁₋₅ branched or unbranched alkyl, phenyl, naphthyl or heterocyclic groups, with each such heterocyclic group being independently selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, isoxazolyl and isothiazolyl, and each such phenyl, naphthyl or heterocyclic group being substituted with 0 to 5 groups selected from halogen, C₁₋₆ branched or unbranched alkyl which is optionally partially or fully halogenated, cyclopropyl, cyclobutyl, cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl and bicycloheptanyl, hydroxy, cyano, C₁₋₃ alkyloxy which is optionally partially or fully halogenated, NH₂C(O), mono- or di(C₁₋₃)alkylaminocarbonyl;
- d) C₅₋₇ cycloalkenyl selected from the group consisting of cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptenyl, cycloheptadienyl, bicyclohexenyl and bicycloheptenyl, wherein such cycloalkenyl group may optionally be substituted with one to three C₁₋₃ alkyl groups;
- e) cyano; and,
- f) methoxycarbonyl, ethoxycarbonyl and propoxycarbonyl;

R₂ is selected from the group consisting of:

a C₁₋₆ branched or unbranched alkyl which may optionally be partially or fully halogenated, acetyl, aroyl, C₁₋₄ branched or unbranched alkoxy, which may optionally be partially or fully halogenated, halogen, methoxycarbonyl and phenylsulfonyl;

R₃ is selected from the group consisting of:

- a) a phenyl, naphthyl or heterocyclic group selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, tetrahydrofuryl, isoxazolyl, isothiazolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, benzoxazolyl, benzisoxazolyl, benzpyrazolyl, benzothiofuranyl, cinnolyl, pterindinyl, phthalazinyl, naphthypyridinyl, quinoxalyl, quinazolinyl, purinyl and indazolyl; wherein such phenyl, naphthyl or heterocyclic group is optionally substituted with one to five groups selected from the group consisting of a C₁₋₆ branched or unbranched alkyl, phenyl, naphthyl, heterocycle selected from the group hereinabove described, C₁₋₆ branched or unbranched alkyl which is optionally partially or fully halogenated, cyclopropyl, cyclobutyl, cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl, bicycloheptanyl, phenyl C₁₋₅ alkyl, naphthyl C₁₋₅ alkyl, halo, hydroxy, cyano, C₁₋₃ alkyloxy which may optionally be partially or fully halogenated, phenyloxy, naphthyloxy, heteraryloxy wherein the heterocyclic moiety is selected from the group hereinabove described, nitro, amino, mono- or di-(C₁₋₃)alkylamino, phenylamino, naphthylamino, heterocyclylamino wherein the heterocyclyl moiety is selected from the group hereinabove described, NH₂C(O), a mono- or di-(C₁₋₃)alkyl aminocarbonyl, C₁₋₅ alkyl-C(O)-C₁₋₄ alkyl, amino-C₁₋₅ alkyl, mono- or di-(C₁₋₃)alkylamino-C₁₋₅ alkyl, amino-S(O)₂, di-(C₁₋₃)alkylamino-S(O)₂, R₄-C₁₋₅ alkyl, R₅-C₁₋₅ alkoxy, R₆-C(O)-C₁₋₅ alkyl and R₇-C₁₋₅ alkyl(R₈)N;
- b) a fused aryl selected from the group consisting of benzocyclobutanyl, indanyl, indenyl, dihydronaphthyl, tetrahydronaphthyl, benzocycloheptanyl and benzocycloheptenyl, or a fused heterocyclyl selected from the group consisting of cyclopentenopyridine, cyclohexanopyridine, cyclopentanopyrimidine, cyclohexanopyrimidine, cyclopentanopyrazine, cyclohexanopyrazine, cyclopentanopyridazine, cyclohexanopyridazine, cyclopentanoquinoline, cyclohexanoquinoline, cyclopentanoisoquinoline, cyclohexanoisoquinoline, cyclopentanoindole, cyclohexanoindole, cyclopentanobenzimidazole, cyclohexanobenzimidazole, cyclopentanobenzoxazole, cyclohexanobenzoxazole, cyclopentanoimidazole, cyclohexanoimidazole, cyclopentanothiophene and cyclohexanothiophene; wherein the fused aryl or fused heterocyclyl ring is substituted with 0 to 3 groups independently selected from phenyl, naphthyl and heterocyclyl selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl,

pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, isoxazolyl, and isothiazolyl, C₁₋₆ branched or unbranched alkyl which is optionally partially or fully halogenated, halo, cyano, C₁₋₃ alkyloxy which is optionally partially or fully halogenated, phenyloxy, naphthyloxy, heterocyclyloxy wherein the heterocyclyl moiety is selected from the group hereinabove described, nitro, amino, mono- or di-(C₁₋₃)alkylamino, phenylamino, naphthylamino, heterocyclylamino wherein the heterocyclyl moiety is selected from the group hereinabove described, NH₂C(O), a mono- or di-(C₁₋₃)alkyl aminocarbonyl, C₁₋₄ alkyl-OC(O), C₁₋₅ alkyl-C(O)-C₁₋₄ branched or unbranched alkyl, an amino-C₁₋₅ alkyl, mono- or di-(C₁₋₃)alkylamino-C₁₋₅ alkyl, R₉-C₁₋₅ alkyl, R₁₀-C₁₋₅ alkoxy, R₁₁-C(O)-C₁₋₅ alkyl and R₁₂-C₁₋₅ alkyl(R₁₃)N;

c) cycloalkyl selected from the group consisting of cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl and bicycloheptanyl, which the cycloalkyl may optionally be partially or fully halogenated and which may optionally be substituted with one to three C₁₋₃ alkyl groups;

d) C₅₋₇ cycloalkenyl, selected from the group consisting of cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptenyl, cycloheptadienyl, bicyclohexenyl and bicycloheptenyl, wherein such cycloalkenyl group may optionally be substituted with one to three C₁₋₃ alkyl groups; and

e) acetyl, aroyl, alkoxy carbonyl alkyl or phenylsulfonyl;

f) C₁₋₆ branched or unbranched alkyl which may optionally be partially or fully halogenated;

wherein

or R₁ and R₂ taken together may optionally form a fused phenyl or pyridinyl ring,

each R₈, R₁₃ is independently selected from the group consisting of:

hydrogen and C₁₋₄ branched or unbranched alkyl which may optionally be partially or fully halogenated;

each R_4 , R_5 , R_6 , R_7 , R_9 , R_{10} , R_{11} and R_{12} is independently selected from the group consisting of:
morpholine, piperidine, piperazine, imidazole and tetrazole;

$m = 0, 1, 2$;

$r = 0, 1, 2$;

$t = 0, 1, 2$;

and


$X = O$ or S or

the physiologically acceptable acids or salts thereof.

Claim 21 (previously presented): The method according to claim 20 wherein the compound is:

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea
or the physiologically acceptable acids or salts thereof.

Claim 22 (currently amended): A method of treating proliferation of acute myelogenous leukemia blasts or plasma cell dyscrasias said method comprising administering to a patient in need of such treatment a therapeutically effective amount of a compound of the formula (I):



wherein

Ar_1 is a pyrazole wherein Ar_1 may be substituted by one or more R_1, R_2 or R_3 ;

Ar₂ is:

phenyl, naphthyl, quinoline, isoquinoline, tetrahydronaphthyl, tetrahydroquinoline, tetrahydroisoquinoline, benzimidazole, benzofuran, indanyl, indenyl or indole each being optionally substituted with one to three R₂ groups;

L is a C₁₋₁₀ saturated or unsaturated branched or unbranched carbon chain;

wherein one or more methylene groups are optionally independently replaced by O, N or S; and

wherein said linking group is optionally substituted with 0-2 oxo groups and one or more C₁₋₄ branched or unbranched alkyl which may be substituted by one or more halogen atoms;

Q is selected from the group consisting of:

- a) pyridine, pyrimidine, pyridazine, imidazole, benzimidazole, oxazo[4,5-*b*]pyridine and imidazo[4,5-*b*]pyridine, which are optionally substituted with one to three groups selected from the group consisting of halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, hydroxy, mono- or di-(C₁₋₃ alkyl)amino, C₁₋₆ alkyl-S(O)_m and phenylamino wherein the phenyl ring is optionally substituted with one to two groups consisting of halogen, C₁₋₆ alkyl and C₁₋₆ alkoxy;
- b) morpholine, thiomorpholine, thiomorpholine sulfoxide, thiomorpholine sulfone, piperidine, piperidinone and tetrahydropyrimidone which are optionally substituted with one to three groups selected from the group consisting of C₁₋₆ alkyl, C₁₋₆ alkoxy, hydroxy, mono- or di-(C₁₋₃ alkyl)amino-C₁₋₃ alkyl, phenylamino-C₁₋₃ alkyl and C₁₋₃ alkoxy-C₁₋₃ alkyl;

R₁ is selected from the group consisting of:

- a) C₃₋₁₀ branched or unbranched alkyl, which may optionally be partially or fully halogenated, and optionally substituted with one to three phenyl, naphthyl or heterocyclic groups selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl,

- pyrazolyl, thienyl, furyl, isoxazolyl and isothiazolyl; each such phenyl, naphthyl or heterocycle selected from the group hereinabove described, being substituted with 0 to 5 groups selected from the group consisting of halogen, C₁₋₆ branched or unbranched alkyl which is optionally partially or fully halogenated, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkenyl, hydroxy, cyano, C₁₋₃ alkyloxy which is optionally partially or fully halogenated, NH₂C(O) and di(C₁₋₃)alkylaminocarbonyl;
- b) C₃₋₇ cycloalkyl selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl and bicycloheptanyl, which may optionally be partially or fully halogenated and which may optionally be substituted with one to three C₁₋₃ alkyl groups, or an analog of such cycloalkyl group wherein one to three ring methylene groups are replaced by groups independently selected from O, S, CHOH, >C=O, >C=S and NH;
- c) C₃₋₁₀ branched alkenyl which may optionally be partially or fully halogenated, and which is optionally substituted with one to three C₁₋₅ branched or unbranched alkyl, phenyl, naphthyl or heterocyclic groups, with each such heterocyclic group being independently selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, isoxazolyl and isothiazolyl, and each such phenyl, naphthyl or heterocyclic group being substituted with 0 to 5 groups selected from halogen, C₁₋₆ branched or unbranched alkyl which is optionally partially or fully halogenated, cyclopropyl, cyclobutyl, cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl and bicycloheptanyl, hydroxy, cyano, C₁₋₃ alkyloxy which is optionally partially or fully halogenated, NH₂C(O), mono- or di(C₁₋₃)alkylaminocarbonyl;
- d) C₅₋₇ cycloalkenyl selected from the group consisting of cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptenyl, cycloheptadienyl, bicyclohexenyl and bicycloheptenyl, wherein such cycloalkenyl group may optionally be substituted with one to three C₁₋₃ alkyl groups;
- e) cyano; and,
- f) methoxycarbonyl, ethoxycarbonyl and propoxycarbonyl;

R₂ is selected from the group consisting of:

a C₁₋₆ branched or unbranched alkyl which may optionally be partially or fully halogenated, acetyl, aroyl, C₁₋₄ branched or unbranched alkoxy, which may optionally be partially or fully halogenated, halogen, methoxycarbonyl and phenylsulfonyl;

R₃ is selected from the group consisting of:

- a) a phenyl, naphthyl or heterocyclic group selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, tetrahydrofuryl, isoxazolyl, isothiazolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, benzoxazolyl, benzisoxazolyl, benzpyrazolyl, benzothiofuranyl, cinnolyl, pterindinyl, phthalazinyl, naphthypyridinyl, quinoxalyl, quinazolinyl, purinyl and indazolyl; wherein such phenyl, naphthyl or heterocyclic group is optionally substituted with one to five groups selected from the group consisting of a C₁₋₆ branched or unbranched alkyl, phenyl, naphthyl, heterocycle selected from the group hereinabove described, C₁₋₆ branched or unbranched alkyl which is optionally partially or fully halogenated, cyclopropyl, cyclobutyl, cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl, bicycloheptanyl, phenyl C₁₋₅ alkyl, naphthyl C₁₋₅ alkyl, halo, hydroxy, cyano, C₁₋₃ alkoxy which may optionally be partially or fully halogenated, phenyloxy, naphthyloxy, heteraryloxy wherein the heterocyclic moiety is selected from the group hereinabove described, nitro, amino, mono- or di-(C₁₋₃)alkylamino, phenylamino, naphthylamino, heterocyclamino wherein the heterocycl moiety is selected from the group hereinabove described, NH₂C(O), a mono- or di-(C₁₋₃)alkyl aminocarbonyl, C₁₋₅ alkyl-C(O)-C₁₋₄ alkyl, amino-C₁₋₅ alkyl, mono- or di-(C₁₋₃)alkylamino-C₁₋₅ alkyl, amino-S(O)₂, di-(C₁₋₃)alkylamino-S(O)₂, R₄-C₁₋₅ alkyl, R₅-C₁₋₅ alkoxy, R₆-C(O)-C₁₋₅ alkyl and R₇-C₁₋₅ alkyl(R₈)N;
- b) a fused aryl selected from the group consisting of benzocyclobutanyl, indanyl, indenyl, dihydronaphthyl, tetrahydronaphthyl, benzocycloheptanyl and benzocycloheptenyl, or a fused heterocycl selected from the group consisting of cyclopentenopyridine, cyclohexanopyridine, cyclopentanopyrimidine, cyclohexanopyrimidine, cyclopentanopyrazine, cyclohexanopyrazine, cyclopentanopyridazine, cyclohexanopyridazine, cyclopentanoquinoline, cyclohexanoquinoline,

cyclopentanoisoquinoline, cyclohexanoisoquinoline, cyclopentanoindole, cyclohexanoindole, cyclopentanobenzimidazole, cyclohexanobenzimidazole, cyclopentanobenzoxazole, cyclohexanobenzoxazole, cyclopentanoimidazole, cyclohexanoimidazole, cyclopentanothiophene and cyclohexanothiophene; wherein the fused aryl or fused heterocyclyl ring is substituted with 0 to 3 groups independently selected from phenyl, naphthyl and heterocyclyl selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, isoxazolyl, and isothiazolyl, C₁₋₆ branched or unbranched alkyl which is optionally partially or fully halogenated, halo, cyano, C₁₋₃ alkyloxy which is optionally partially or fully halogenated, phenyloxy, naphthyloxy, heterocyclyloxy wherein the heterocyclyl moiety is selected from the group hereinabove described, nitro, amino, mono- or di-(C₁₋₃)alkylamino, phenylamino, naphthylamino, heterocyclylamino wherein the heterocyclyl moiety is selected from the group hereinabove described, NH₂C(O), a mono- or di-(C₁₋₃)alkyl aminocarbonyl, C₁₋₄ alkyl-OC(O), C₁₋₅ alkyl-C(O)-C₁₋₄ branched or unbranched alkyl, an amino-C₁₋₅ alkyl, mono- or di-(C₁₋₃)alkylamino-C₁₋₅ alkyl, R₉-C₁₋₅ alkyl, R₁₀-C₁₋₅ alkoxy, R₁₁-C(O)-C₁₋₅ alkyl and R₁₂-C₁₋₅ alkyl(R₁₃)N;

c) cycloalkyl selected from the group consisting of cyclopentanyl, cyclohexanyl, cycloheptanyl, bicyclopentanyl, bicyclohexanyl and bicycloheptanyl, which the cycloalkyl may optionally be partially or fully halogenated and which may optionally be substituted with one to three C₁₋₃ alkyl groups;

d) C₅₋₇ cycloalkenyl, selected from the group consisting of cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptenyl, cycloheptadienyl, bicyclohexenyl and bicycloheptenyl, wherein such cycloalkenyl group may optionally be substituted with one to three C₁₋₃ alkyl groups; and

e) acetyl, aroyl, alkoxycarbonylalkyl or phenylsulfonyl;

f) C₁₋₆ branched or unbranched alkyl which may optionally be partially or fully halogenated;

wherein

or R₁ and R₂ taken together may optionally form a fused phenyl or pyridinyl ring,

each R₈, R₁₃ is independently selected from the group consisting of:

hydrogen and C₁₋₄ branched or unbranched alkyl which may optionally be partially or fully halogenated;

each R₄, R₅, R₆, R₇, R₉, R₁₀, R₁₁ and R₁₂ is independently selected from the group consisting of:
morpholine, piperidine, piperazine, imidazole and tetrazole;

m = 0, 1, 2;

r = 0, 1, 2;

t = 0, 1, 2;

and

X = O or S or

the physiologically acceptable acids or salts thereof.

Claim 23 (previously presented): The method according to claim 22 wherein the compound is:

1-[5-*tert*-Butyl-2-*p*-tolyl-2H-pyrazol-3-yl]-3-[4-(2-morpholin-4-yl-ethoxy)naphthalen-1-yl]-urea
or the physiologically acceptable acids or salts thereof.